Web application for Intensity of Erosion and Outflow

Name of the River Basin: Shirindareh S10-1

Country: Iran, Islamic Republic of

Year: 2019

GPS coordinates, latitude and longitude with Google Maps: 37.81,57.19

INPUT DATA

Geometric characteristics of the river basins

F = 15.92 km² (Surface area of the drainage basin)

O = 17.41 km (Length of the watershed)

 $Fv = 8.46 \text{ km}^2$ (Surface area of greater portion of the drainage basin)

Fm = 7.46 km² (Surface area of smaller portion of the drainage basin)

Lv = 7.3 km (Natural length of main water course)

Lb = 7.18 km (Length of the drainage basin measured by a series of paraller lines)

Topograpfic characteristics of the river basins

Contour line length - Liz [km]: ["2.33 ","9.15 ","13.95 ","15.71 ","13.94 ","10.72 ","4.53 ","0.99 "]

The area between the two neighboring contour lines - f [km²]: ["0.34 ","1.69 ","2.57 ","3.50 ","3.11 ","3.01 ","1.43 ","0.26 ","0.01 "]

h0 = 900 m (Altitude of the initial contour)

 $\Delta h = 100 \text{ m (Equidistance)}$

Hmin = 831 (Lowest altitude in the drainage basin)

Hmax = 1634 (Highest altitude in the draigane basin

Hydrological characteristics of the river basins

 $\Sigma L = 24.83$ km (The total length of the main watercourse with tributaries of 1st and 2nd class)

Lm = 6.46 km (The shortest distance between the fountain (head and mouth))

Water permeability

fp = 0.03 (Part of the surface area of the drainage basin which is composed of highly water permeable structures from the rocks (limestone, sand, gravel))

fpp = 0 (Part of the surface area of the drainage basin which is composed of the rocks of medium water permeability (schist, marls, sandstone))

fo = 0.97 (Part of the surface area of the drainage basin which is composed of the rocks of poor water permeability (heavy clay, compact eruptive))

Land use

fs = 0 (Part of the surface area of the drainage basin under the forest)

ft = 1.00000 (Part of the surface area of the drainage basin which is under the grass, meadows, pastures and orchards)

fg = 0.00000 (Part of the surface area of the drainage basin which is bare or under the soils without grass vegetation)

Meteorological data

hb = 32.76 mm (Level of torrent rain)

Up (years) = 100

to = 12.40 °C (Average annual air temperature)

Hgod = 293.9 mm (Average annual quantity of precipitation)

Erosion coefficients

Y = 1.1793 (Types of soil structures and allied types)

0 % (Sand, gravel and incoherent soils)

0 % (Saline soils)

96.55 % (Decomposed limestone and marls)

0 % (Serpentines, red sand stones, flishe deposits)

0 % (Podzols and parapodzols, decomposed schist)

0 % (Solid and Schist limestone, Terra Rosa and Humic soil)

0 % (Brown forest soils and Mountain soils)

```
3.45 % (Epieugleysol and Marshlands)
0 % (Good structured Chernozems and alluvial well-structured deposits)
0 % (Bare, compact igneous)
Xa = 0.7 (Planning of the drainage basin, rate of drainage basin regulation)
0 % (Bare lands)
0 % (Plough-lands)
100 % (Orchards and vineyards)
0 % (Mountain pastures)
0 % (Meadows)
0 % (Degraded forests)
0 % (Well-constituted forests)
\phi = 0.32941 (Numerical coefficient of visible and clearly pointed processes of soil erosion)
0 % (Depth erosion)
3.56 % (80% of the river basin under rill and gully erosion)
1.61 % (50% of the river basin under rill and gully erosion)
0 % (100% of the river basin under surface erosion)
0 % (100% of the river basin under surface erosion, without visible furrows, ravines and land
slides)
0 % (50% of the river basin under surface erosion)
94.83 % (20% of the river basin under surface erosion)
0 % (There are smaller slides in the watercourse beds)
0 % (The river basin mostly under plough-land)
0 % (The river basin under forests and perennial vegetation)
INPUT DATA
A = 0.46506164383562 (Coefficient of the river basin form)
```

m = 0.51611489913903 (Coefficient of the watershed development) **B** = 2.2172701949861 km (Average river basin width) a = 0.12562814070352 ((A)symmetry of the river basin)

G = 1.5596733668342 (Density of the river network of the basin)

K = 1.1300309597523 (Coefficient of the river basin tortuousness)

 $H_{sr} = 1197.6721105528 \text{ m}$ (Average river basin altitude)

D=366.6721105528 m (Average elevation difference of the river basin) $I_{\rm sr}=44.798994974874$ % (Average river basin decline) $H_{\rm leb}=803$ m (The height of the local erosion base of the river basin)

 $E_r = 127.96167218094$ (Coefficient of the erosion energy of the river basins relief)

 $S_1 = 0.982$ (Coefficient of the regions permeability)

 $S_2 = 0.8$ (Coefficient of the vegetation cover)

W = 0.44123541903715 m (Analytical presentation of the water retention in inflow)

 $2gDF^{1/2} = 338.42307900025 \text{ m km s}^{-1}$ (Energetic potential of water flow during torrent rains)

 $Q_{max} = 54.555976868649 \text{ m}^3 \text{ s}^{-1}$ (Maximal outflow from the river basin)

T = 1.157583690279 (Temperature coefficient of the region)

Z = 0.82446202617277 (Coefficient of the river basin erosion)

 $W_{god} = 12737.990108741 \text{ m}^3 \text{ god}^{-1}$ (Production of erosion material in the river basin

 $R_u = 0.29209371831941$ (Coefficient of the deposit retention)

 $G_{god} = 3720.686894778 \text{ m}^3 \text{ god}^{-1} \text{ (Real soil losses)}$

 $G_{god} \text{ km}^{-2} = 233.71148836545 \text{ m}^3 \text{ km}^{-2} \text{ god}^{-1} \text{ (Real soil losses per km}^2\text{)}$

http://www.wintero.me