
Web application for Intensity of Erosion and Outflow

Name of the River Basin: Shirindareh S1-intB

Country: Iran, Islamic Republic of

Year: 2019

GPS coordinates, latitude and longitude with Google Maps: 37.73,57.67

INPUT DATA

Geometric characteristics of the river basins

F = 69.37 km² (Surface area of the drainage basin)

O = 57.79 km (Length of the watershed)

Fv = 46.42 km² (Surface area of greater portion of the drainage basin)

Fm = 22.95 km² (Surface area of smaller portion of the drainage basin)

Lv = 13.05 km (Natural length of main water course)

Lb = 16.05 km (Length of the drainage basin measured by a series of parallel lines)

Topographic characteristics of the river basins

Contour line length - Liz [km]: ["16.18 ", "59.74 ", "45.02 ", "28.93 ", "24.26 ", "8.28 ", "0.17 "]

The area between the two neighboring contour lines - f [km²]: ["3.54 ", "17.82 ", "19.89 ", "11.24 ", "10.91 ", "5.38 ", "0.57 ", "0.02 "]

h0 = 1300 m (Altitude of the initial contour)

Δh = 100 m (Equidistance)

Hmin = 1250 (Lowest altitude in the drainage basin)

Hmax = 1910 (Highest altitude in the drainage basin)

Hydrological characteristics of the river basins

$\Sigma L = 137.92$ km (The total length of the main watercourse with tributaries of 1st and 2nd class)

$L_m = 7.09$ km (The shortest distance between the fountain (head and mouth))

Water permeability

$f_p = 0.17$ (Part of the surface area of the drainage basin which is composed of highly water permeable structures from the rocks (limestone, sand, gravel))

$f_{pp} = 0.38$ (Part of the surface area of the drainage basin which is composed of the rocks of medium water permeability (schist, marls, sandstone))

$f_o = 0.45$ (Part of the surface area of the drainage basin which is composed of the rocks of poor water permeability (heavy clay, compact eruptive))

Land use

$f_s = 0.00$ (Part of the surface area of the drainage basin under the forest)

$f_t = 0.41210$ (Part of the surface area of the drainage basin which is under the grass, meadows, pastures and orchards)

$f_g = 0.58790$ (Part of the surface area of the drainage basin which is bare or under the soils without grass vegetation)

Meteorological data

$h_b = 34.74$ mm (Level of torrent rain)

U_p (years) = 100

$t_o = 10.9$ °C (Average annual air temperature)

$H_{god} = 317.7$ mm (Average annual quantity of precipitation)

Erosion coefficients

$Y = 1.07663$ (Types of soil structures and allied types)

0 % (Sand, gravel and incoherent soils)

0 % (Saline soils)

60.84 % (Decomposed limestone and marls)

22.11 % (Serpentines, red sand stones, flishe deposits)

0.26 % (Podzols and parapodzols, decomposed schist)

0 % (Solid and Schist limestone, Terra Rosa and Humic soil)

0 % (Brown forest soils and Mountain soils)

16.79 % (Epieugleysol and Marshlands)

0 % (Good structured Chernozems and alluvial well-structured deposits)

0 % (Bare, compact igneous)

Xa = 0.77714 (Planning of the drainage basin, rate of drainage basin regulation)

0 % (Bare lands)

58.79 % (Plough-lands)

0.77 % (Orchards and vineyards)

40.44 % (Mountain pastures)

0 % (Meadows)

0 % (Degraded forests)

0 % (Well-constituted forests)

$\phi = 0.66648$ (Numerical coefficient of visible and clearly pointed processes of soil erosion)

0.31 % (Depth erosion)

40.63 % (80% of the river basin under rill and gully erosion)

23.67 % (50% of the river basin under rill and gully erosion)

0 % (100% of the river basin under surface erosion)

0.78 % (100% of the river basin under surface erosion, without visible furrows, ravines and land slides)

0 % (50% of the river basin under surface erosion)

34.45 % (20% of the river basin under surface erosion)

0.16 % (There are smaller slides in the watercourse beds)

0 % (The river basin mostly under plough-land)

0 % (The river basin under forests and perennial vegetation)

INPUT DATA

A = 0.86352873563218 (Coefficient of the river basin form)

m = 0.44199741806007 (Coefficient of the watershed development)

B = 4.3221183800623 km (Average river basin width)

a = 0.67666138100043 ((A)symmetry of the river basin)

G = 1.9881793282399 (Density of the river network of the basin)

K = 1.8406205923836 (Coefficient of the river basin tortuousness)

H_{sr} = 1489.7232232954 m (Average river basin altitude)

D = 239.7232232954 m (Average elevation difference of the river basin)

I_{sr} = 26.319734755658 % (Average river basin decline)

H_{leb} = 660 m (The height of the local erosion base of the river basin)

E_r = 72.79493108475 (Coefficient of the erosion energy of the river basins relief)

S₁ = 0.784 (Coefficient of the regions permeability)

S₂ = 0.91758 (Coefficient of the vegetation cover)

W = 0.45689966318995 m (Analytical presentation of the water retention in inflow)

2gDF^{1/2} = 571.20289915234 m km s⁻¹ (Energetic potential of water flow during torrent rains)

Q_{max} = 162.12427116266 m³ s⁻¹ (Maximal outflow from the river basin)

T = 1.0908712114636 (Temperature coefficient of the region)

Z = 0.9868848776247 (Coefficient of the river basin erosion)

W_{god} = 74047.773412435 m³ god⁻¹ (Production of erosion material in the river basin)

R_u = 0.32295385090679 (Coefficient of the deposit retention)

G_{god} = 23914.01357462 m³ god⁻¹ (Real soil losses)

G_{god} km⁻² = 344.73134747902 m³ km⁻² god⁻¹ (Real soil losses per km²)

<http://www.wintero.me>