# Web application for Intensity of Erosion and Outflow

Name of the River Basin: Shirindareh S6-int

Country: Iran, Islamic Republic of

Year: 2019

GPS coordinates, latitude and longitude with Google Maps: 37.79,57.33

#### **INPUT DATA**

#### Geometric characteristics of the river basins

 $F = 47.66 \text{ km}^2$  (Surface area of the drainage basin)

O = 38.9 km (Length of the watershed)

 $Fv = 35.88 \text{ km}^2$  (Surface area of greater portion of the drainage basin)

Fm = 11.78 km<sup>2</sup> (Surface area of smaller portion of the drainage basin)

Lv = 9.45 km (Natural length of main water course)

Lb = 12.23 km (Length of the drainage basin measured by a series of paraller lines)

#### Topograpfic characteristics of the river basins

Contour line length - Liz [km]: ["12.97 ","39.67 ","42.09 ","26.12 ","8.01 ","4.10 ","0.79 "]

The area between the two neighboring contour lines - f [km<sup>2</sup>]: ["3.07 ","16.07 ","12.62 ","9.71 ","3.96 ","1.41 ","0.79 ","0.03 "]

h0 = 1000 m (Altitude of the initial contour)

Ah = 100 m (Equidistance)

**Hmin = 959 (Lowest altitude in the drainage basin)** 

Hmax = 1615 (Highest altitude in the draigane basin

### Hydrological characteristics of the river basins

 $\Sigma L = 82.88$  km (The total length of the main watercourse with tributaries of 1<sup>st</sup> and 2<sup>nd</sup> class)

Lm = 7.62 km (The shortest distance between the fountain (head and mouth))

#### **Water permeability**

fp = 0.16 (Part of the surface area of the drainage basin which is composed of highly water permeable structures from the rocks (limestone, sand, gravel))

fpp = 0.11 (Part of the surface area of the drainage basin which is composed of the rocks of medium water permeability (schist, marls, sandstone))

fo = 0.73 (Part of the surface area of the drainage basin which is composed of the rocks of poor water permeability (heavy clay, compact eruptive))

#### Land use

fs = 0 (Part of the surface area of the drainage basin under the forest)

ft = 0.00000 (Part of the surface area of the drainage basin which is under the grass, meadows, pastures and orchards)

fg = 1.00000 (Part of the surface area of the drainage basin which is bare or under the soils without grass vegetation)

## Meteorological data

hb = 32.42 mm (Level of torrent rain)

Up (years) = 100

to = 12.60 °C (Average annual air temperature)

Hgod = 289.8 mm (Average annual quantity of precipitation)

### **Erosion coefficients**

Y = 1.0703 (Types of soil structures and allied types)

0 % (Sand, gravel and incoherent soils)

0 % (Saline soils)

50.45 % (Decomposed limestone and marls)

33.52 % (Serpentines, red sand stones, flishe deposits)

0 % (Podzols and parapodzols, decomposed schist)

0 % (Solid and Schist limestone, Terra Rosa and Humic soil)

0 % (Brown forest soils and Mountain soils)

```
16.03 % (Epieugleysol and Marshlands)
0 % (Good structured Chernozems and alluvial well-structured deposits)
0 % (Bare, compact igneous)
Xa = 0.9 (Planning of the drainage basin, rate of drainage basin regulation)
0 % (Bare lands)
100 % (Plough-lands)
0 % (Orchards and vineyards)
0 % (Mountain pastures)
0 % (Meadows)
0 % (Degraded forests)
0 % (Well-constituted forests)
\phi = 0.49584 (Numerical coefficient of visible and clearly pointed processes of soil erosion)
0 % (Depth erosion)
11.95 % (80% of the river basin under rill and gully erosion)
13.77 % (50% of the river basin under rill and gully erosion)
0 % (100% of the river basin under surface erosion)
18.43 % (100% of the river basin under surface erosion, without visible furrows, ravines and
land slides)
0 % (50% of the river basin under surface erosion)
55.85 % (20% of the river basin under surface erosion)
0 % (There are smaller slides in the watercourse beds)
0 % (The river basin mostly under plough-land)
0 % (The river basin under forests and perennial vegetation)
INPUT DATA
```

A = 0.80269841269841 (Coefficient of the river basin form)

m = 0.38614450363893 (Coefficient of the watershed development)

**B** = 3.8969746524939 km (Average river basin width)

a = 1.0113302559799 ((A)symmetry of the river basin)

G = 1.7389844733529 (Density of the river network of the basin)

**K** = 1.240157480315 (Coefficient of the river basin tortuousness)

 $H_{sr} = 1158.0841376416$  m (Average river basin altitude)

D = 199.0841376416 m (Average elevation difference of the river basin)  $I_{sr} = 28.063365505665 \% \text{ (Average river basin decline)}$   $H_{leb} = 656 \text{ m (The height of the local erosion base of the river basin)}$   $E_r = 79.472240547171 \text{ (Coefficient of the erosion energy of the river basins relief)}$   $S_1 = 0.871 \text{ (Coefficient of the regions permeability)}$   $S_2 = 1 \text{ (Coefficient of the vegetation cover)}$  W = 0.43327820625508 m (Analytical presentation of the water retention in inflow)  $2gDF^{1/2} = 431.4642824151 \text{ m km s}^{-1} \text{ (Energetic potential of water flow during torrent rains)}$   $Q_{max} = 130.70200613109 \text{ m}^3 \text{ s}^{-1} \text{ (Maximal outflow from the river basin)}$  T = 1.1661903789691 (Temperature coefficient of the region) Z = 0.9879187996873 (Coefficient of the river basin erosion)  $W_{god} = 49688.240803379 \text{ m}^3 \text{ god}^{-1} \text{ (Production of erosion material in the river basin}$ 

 $R_u = 0.28615643708464$  (Coefficient of the deposit retention)

 $G_{god} = 14218.609953299 \text{ m}^3 \text{ god}^{-1} \text{ (Real soil losses)}$ 

 $G_{god}$  km<sup>-2</sup> = 298.33424157152 m<sup>3</sup> km<sup>-2</sup> god<sup>-1</sup> (Real soil losses per km<sup>2</sup>)

http://www.wintero.me