# Web application for Intensity of Erosion and Outflow

Name of the River Basin: Shirindareh S7-intB

**Country: Iran, Islamic Republic of** 

Year: 2019

GPS coordinates, latitude and longitude with Google Maps: 37.88,57.41

#### **INPUT DATA**

### Geometric characteristics of the river basins

F = 4.12 km<sup>2</sup> (Surface area of the drainage basin)

O = 49.06 km (Length of the watershed)

 $Fv = 3.21 \text{ km}^2$  (Surface area of greater portion of the drainage basin)

Fm = 0.91 km<sup>2</sup> (Surface area of smaller portion of the drainage basin)

Lv = 1.47 km (Natural length of main water course)

Lb = 12.89 km (Length of the drainage basin measured by a series of paraller lines)

#### **Topograpfic characteristics of the river basins**

Contour line length - Liz [km]: ["2.46 ","5.75 ","1.68 ","0.12 "]

The area between the two neighboring contour lines - f [km<sup>2</sup>]: ["0.53 ","2.43 ","0.93 ","0.22 ","0.01 "]

h0 = 1200 m (Altitude of the initial contour)

Ah = 100 m (Equidistance)

**Hmin = 1171 (Lowest altitude in the drainage basin)** 

Hmax = 1502 (Highest altitude in the draigane basin

# Hydrological characteristics of the river basins

 $\Sigma L = 8.56$  km (The total length of the main watercourse with tributaries of 1<sup>st</sup> and 2<sup>nd</sup> class)

Lm = 1.3 km (The shortest distance between the fountain (head and mouth))

# Water permeability

fp = 0.14 (Part of the surface area of the drainage basin which is composed of highly water permeable structures from the rocks (limestone, sand, gravel))

fpp = 0.25 (Part of the surface area of the drainage basin which is composed of the rocks of medium water permeability (schist, marls, sandstone))

fo = 0.61 (Part of the surface area of the drainage basin which is composed of the rocks of poor water permeability (heavy clay, compact eruptive))

#### Land use

fs = 0 (Part of the surface area of the drainage basin under the forest)

ft = 0.89070 (Part of the surface area of the drainage basin which is under the grass, meadows, pastures and orchards)

fg = 0.10930 (Part of the surface area of the drainage basin which is bare or under the soils without grass vegetation)

# Meteorological data

hb = 33.2 mm (Level of torrent rain)

Up (years) = 100

to = 12 °C (Average annual air temperature)

Hgod = 299.2 mm (Average annual quantity of precipitation)

# **Erosion coefficients**

**Y = 0.99021 (Types of soil structures and allied types)** 

0 % (Sand, gravel and incoherent soils)

0 % (Saline soils)

7.41 % (Decomposed limestone and marls)

69.15 % (Serpentines, red sand stones, flishe deposits)

0 % (Podzols and parapodzols, decomposed schist)

0 % (Solid and Schist limestone, Terra Rosa and Humic soil)

0 % (Brown forest soils and Mountain soils)

```
23.44 % (Epieugleysol and Marshlands)
0 % (Good structured Chernozems and alluvial well-structured deposits)
0 % (Bare, compact igneous)
Xa = 0.65974 (Planning of the drainage basin, rate of drainage basin regulation)
0 % (Bare lands)
10.93 % (Plough-lands)
26.95 % (Orchards and vineyards)
62.12 % (Mountain pastures)
0 % (Meadows)
0 % (Degraded forests)
0 % (Well-constituted forests)
\phi = 0.4668 (Numerical coefficient of visible and clearly pointed processes of soil erosion)
0 % (Depth erosion)
0 % (80% of the river basin under rill and gully erosion)
33.36 % (50% of the river basin under rill and gully erosion)
0 % (100% of the river basin under surface erosion)
0 % (100% of the river basin under surface erosion, without visible furrows, ravines and land
slides)
0 % (50% of the river basin under surface erosion)
66.64 % (20% of the river basin under surface erosion)
0 % (There are smaller slides in the watercourse beds)
0 % (The river basin mostly under plough-land)
0 % (The river basin under forests and perennial vegetation)
INPUT DATA
A = 6.5079591836735 (Coefficient of the river basin form)
```

m = 0.20429784930106 (Coefficient of the watershed development)

**B** = 0.31962761830877 km (Average river basin width)

a = 1.1165048543689 ((A)symmetry of the river basin)

**G** = 2.0776699029126 (Density of the river network of the basin)

**K** = 1.1307692307692 (Coefficient of the river basin tortuousness)

 $H_{sr} = 1275.5643203883 \text{ m}$  (Average river basin altitude)

D = 104.5643203883 m (Average elevation difference of the river basin)  $I_{sr} = 24.296116504854 \% \text{ (Average river basin decline)}$   $H_{leb} = 331 \text{ m (The height of the local erosion base of the river basin)}$   $E_r = 73.952662952726 \text{ (Coefficient of the erosion energy of the river basins relief)}$   $S_1 = 0.841 \text{ (Coefficient of the regions permeability)}$   $S_2 = 0.82186 \text{ (Coefficient of the vegetation cover)}$  W = 0.46167486176963 m (Analytical presentation of the water retention in inflow)  $2gDF^{1/2} = 91.936902819249 \text{ m km s}^{-1} \text{ (Energetic potential of water flow during torrent rains)}$   $Q_{max} = 190.92586066825 \text{ m}^3 \text{ s}^{-1} \text{ (Maximal outflow from the river basin)}$  T = 1.1401754250991 (Temperature coefficient of the region) Z = 0.62696104243928 (Coefficient of the river basin erosion)  $W_{god} = 2192.0032203827 \text{ m}^3 \text{ god}^{-1} \text{ (Production of erosion material in the river basin)}$   $R_u = 0.39493180420294 \text{ (Coefficient of the deposit retention)}$ 

 $G_{god} = 865.69178664438 \text{ m}^3 \text{ god}^{-1} \text{ (Real soil losses)}$ 

 $G_{god} \text{ km}^{-2} = 210.11936569038 \text{ m}^3 \text{ km}^{-2} \text{ god}^{-1} \text{ (Real soil losses per km}^2\text{)}$ 

http://www.wintero.me