
Web application for Intensity of Erosion and Outflow

Name of the River Basin: Shirindareh S8-intB

Country: Iran, Islamic Republic of

Year: 2019

GPS coordinates, latitude and longitude with Google Maps: 37.87,57.29

INPUT DATA

Geometric characteristics of the river basins

F = 23.48 km² (Surface area of the drainage basin)

O = 24.55 km (Length of the watershed)

Fv = 16.76 km² (Surface area of greater portion of the drainage basin)

Fm = 6.71 km² (Surface area of smaller portion of the drainage basin)

Lv = 4.39 km (Natural length of main water course)

Lb = 12.51 km (Length of the drainage basin measured by a series of parallel lines)

Topographic characteristics of the river basins

Contour line length - Liz [km]: ["14.18 ", "25.53 ", "22.13 ", "5.68 "]

The area between the two neighboring contour lines - f [km²]: ["2.90 ", "7.70 ", "8.92 ", "3.95 ", "0.01 "]

h0 = 1100 m (Altitude of the initial contour)

Δh = 100 m (Equidistance)

Hmin = 1029 (Lowest altitude in the drainage basin)

Hmax = 1494 (Highest altitude in the drainage basin)

Hydrological characteristics of the river basins

$\Sigma L = 42.13$ km (The total length of the main watercourse with tributaries of 1st and 2nd class)

$L_m = 4.24$ km (The shortest distance between the fountain (head and mouth))

Water permeability

$f_p = 0.36$ (Part of the surface area of the drainage basin which is composed of highly water permeable structures from the rocks (limestone, sand, gravel))

$f_{pp} = 0$ (Part of the surface area of the drainage basin which is composed of the rocks of medium water permeability (schist, marls, sandstone))

$f_o = 0.64$ (Part of the surface area of the drainage basin which is composed of the rocks of poor water permeability (heavy clay, compact eruptive))

Land use

$f_s = 0$ (Part of the surface area of the drainage basin under the forest)

$f_t = 0.84750$ (Part of the surface area of the drainage basin which is under the grass, meadows, pastures and orchards)

$f_g = 0.15250$ (Part of the surface area of the drainage basin which is bare or under the soils without grass vegetation)

Meteorological data

$h_b = 32.83$ mm (Level of torrent rain)

U_p (years) = 100

$t_o = 12.30$ °C (Average annual air temperature)

$H_{god} = 294.7$ mm (Average annual quantity of precipitation)

Erosion coefficients

$Y = 1.04975$ (Types of soil structures and allied types)

0 % (Sand, gravel and incoherent soils)

0 % (Saline soils)

0 % (Decomposed limestone and marls)

89.95 % (Serpentines, red sand stones, flishe deposits)

0 % (Podzols and parapodzols, decomposed schist)

0 % (Solid and Schist limestone, Terra Rosa and Humic soil)

0 % (Brown forest soils and Mountain soils)

10.05 % (Epieugleysol and Marshlands)

0 % (Good structured Chernozems and alluvial well-structured deposits)

0 % (Bare, compact igneous)

$X_a = 0.64575$ (Planning of the drainage basin, rate of drainage basin regulation)

0 % (Bare lands)

15.25 % (Plough-lands)

0 % (Orchards and vineyards)

84.75 % (Mountain pastures)

0 % (Meadows)

0 % (Degraded forests)

0 % (Well-constituted forests)

$\phi = 0.45258$ (Numerical coefficient of visible and clearly pointed processes of soil erosion)

0 % (Depth erosion)

8.48 % (80% of the river basin under rill and gully erosion)

20.34 % (50% of the river basin under rill and gully erosion)

0 % (100% of the river basin under surface erosion)

0 % (100% of the river basin under surface erosion, without visible furrows, ravines and land slides)

0 % (50% of the river basin under surface erosion)

71.18 % (20% of the river basin under surface erosion)

0 % (There are smaller slides in the watercourse beds)

0 % (The river basin mostly under plough-land)

0 % (The river basin under forests and perennial vegetation)

INPUT DATA

$A = 1.0904897494305$ (Coefficient of the river basin form)

$m = 0.25557039367567$ (Coefficient of the watershed development)

$B = 1.876898481215$ km (Average river basin width)

$a = 0.85641244141457$ ((A)symmetry of the river basin)

$G = 1.7942930153322$ (Density of the river network of the basin)

$K = 1.0353773584906$ (Coefficient of the river basin tortuousness)

$H_{sr} = 1211.2018739353$ m (Average river basin altitude)

$D = 182.2018739353$ m (Average elevation difference of the river basin)

$I_{sr} = 28.756388415673$ % (Average river basin decline)

$H_{leb} = 465$ m (The height of the local erosion base of the river basin)

$E_r = 67.240131955776$ (Coefficient of the erosion energy of the river basins relief)

$S_1 = 0.784$ (Coefficient of the regions permeability)

$S_2 = 0.8305$ (Coefficient of the vegetation cover)

$W = 0.44810225749552$ m (Analytical presentation of the water retention in inflow)

$2gDF^{1/2} = 289.71765910972$ m km s⁻¹ (Energetic potential of water flow during torrent rains)

$Q_{max} = 92.178446878278$ m³ s⁻¹ (Maximal outflow from the river basin)

$T = 1.1532562594671$ (Temperature coefficient of the region)

$Z = 0.67030407415959$ (Coefficient of the river basin erosion)

$W_{god} = 13758.202908598$ m³ god⁻¹ (Production of erosion material in the river basin)

$R_u = 0.2939485820865$ (Coefficient of the deposit retention)

$G_{god} = 4044.2042370407$ m³ god⁻¹ (Real soil losses)

$G_{god} \text{ km}^{-2} = 172.24038488248$ m³ km⁻² god⁻¹ (Real soil losses per km²)

<http://www.wintero.me>