Web application for Intensity of Erosion and Outflow

Name of the River Basin: Shirindareh S10-intC

Country: Iran, Islamic Republic of

Year: 2019

GPS coordinates, latitude and longitude with Google Maps: 37.74,57.17

INPUT DATA

Geometric characteristics of the river basins

F = 2.31 km² (Surface area of the drainage basin)

O = 8.08 km (Length of the watershed)

 $Fv = 1.95 \text{ km}^2$ (Surface area of greater portion of the drainage basin)

Fm = 0.36 km² (Surface area of smaller portion of the drainage basin)

Lv = 1.66 km (Natural length of main water course)

Lb = 9.11 km (Length of the drainage basin measured by a series of paraller lines)

Topograpfic characteristics of the river basins

Contour line length - Liz [km]: ["2.73 ","0.60 "]

The area between the two neighboring contour lines - f [km²]: ["1.67 ","0.63 ","0.01 "]

h0 = 800 m (Altitude of the initial contour)

 $\Delta h = 100 \text{ m (Equidistance)}$

Hmin = 742 (Lowest altitude in the drainage basin)

Hmax = 925 (Highest altitude in the draigane basin

Hydrological characteristics of the river basins

 $\Sigma L = 1.97$ km (The total length of the main watercourse with tributaries of 1st and 2nd class)

Lm = 1.3 km (The shortest distance between the fountain (head and mouth))

Water permeability

fp = 0.42 (Part of the surface area of the drainage basin which is composed of highly water permeable structures from the rocks (limestone, sand, gravel))

fpp = 0.58 (Part of the surface area of the drainage basin which is composed of the rocks of medium water permeability (schist, marls, sandstone))

fo = 0 (Part of the surface area of the drainage basin which is composed of the rocks of poor water permeability (heavy clay, compact eruptive))

Land use

fs = 0 (Part of the surface area of the drainage basin under the forest)

ft = 0.81090 (Part of the surface area of the drainage basin which is under the grass, meadows, pastures and orchards)

fg = 0.18910 (Part of the surface area of the drainage basin which is bare or under the soils without grass vegetation)

Meteorological data

hb = 29.87 mm (Level of torrent rain)

Up (years) = 100

to = 14.60 °C (Average annual air temperature)

Hgod = 258.9 mm (Average annual quantity of precipitation)

Erosion coefficients

Y = 0.89355 (Types of soil structures and allied types)

0 % (Sand, gravel and incoherent soils)

0 % (Saline soils)

0 % (Decomposed limestone and marls)

58.71 % (Serpentines, red sand stones, flishe deposits)

0 % (Podzols and parapodzols, decomposed schist)

0 % (Solid and Schist limestone, Terra Rosa and Humic soil)

0 % (Brown forest soils and Mountain soils)

```
41.29 % (Epieugleysol and Marshlands)
0 % (Good structured Chernozems and alluvial well-structured deposits)
0 % (Bare, compact igneous)
Xa = 0.6733 (Planning of the drainage basin, rate of drainage basin regulation)
1.99 % (Bare lands)
16.92 % (Plough-lands)
14.58 % (Orchards and vineyards)
66.51 % (Mountain pastures)
0 % (Meadows)
0 % (Degraded forests)
0 % (Well-constituted forests)
\phi = 0.79338 (Numerical coefficient of visible and clearly pointed processes of soil erosion)
0 % (Depth erosion)
17.98 % (80% of the river basin under rill and gully erosion)
77.1 % (50% of the river basin under rill and gully erosion)
0 % (100% of the river basin under surface erosion)
0 % (100% of the river basin under surface erosion, without visible furrows, ravines and land
slides)
0 % (50% of the river basin under surface erosion)
4.92 % (20% of the river basin under surface erosion)
0 % (There are smaller slides in the watercourse beds)
0 % (The river basin mostly under plough-land)
0 % (The river basin under forests and perennial vegetation)
INPUT DATA
```

A = 0.94915662650602 (Coefficient of the river basin form) m = 0.30810387900233 (Coefficient of the watershed development) B = 0.25356750823271 km (Average river basin width) a = 1.3766233766234 ((A)symmetry of the river basin) G = 0.85281385281385 (Density of the river network of the basin) **K = 1.2769230769231 (Coefficient of the river basin tortuousness)**

 $H_{sr} = 793.15800865801 \text{ m}$ (Average river basin altitude)

D = 51.15800865801 m (Average elevation difference of the river basin) $I_{sr} = 14.415584415584$ % (Average river basin decline)

 H_{leb} = 183 m (The height of the local erosion base of the river basin)

 $E_r = 47.24960933478$ (Coefficient of the erosion energy of the river basins relief)

 $S_1 = 0.574$ (Coefficient of the regions permeability)

 $S_2 = 0.83782$ (Coefficient of the vegetation cover)

W = 0.41687578733098 m (Analytical presentation of the water retention in inflow)

 $2gDF^{1/2} = 48.151775668194 \text{ m km s}^{-1}$ (Energetic potential of water flow during torrent rains)

 $Q_{max} = 9.1626158413658 \text{ m}^3 \text{ s}^{-1}$ (Maximal outflow from the river basin)

T = 1.2489995996797 (Temperature coefficient of the region)

Z = 0.70574398239862 (Coefficient of the river basin erosion)

 $W_{god} = 1391.3198390988 \text{ m}^3 \text{ god}^{-1}$ (Production of erosion material in the river basin

 $R_u = 0.11027928487097$ (Coefficient of the deposit retention)

 $G_{god} = 153.4337568826 \text{ m}^3 \text{ god}^{-1} \text{ (Real soil losses)}$

 G_{god} km⁻² = 66.421539776019 m³ km⁻² god⁻¹ (Real soil losses per km²)

http://www.wintero.me