Web application for Intensity of Erosion and Outflow

Name of the River Basin: Shirindareh S2-intA

Country: Iran, Islamic Republic of

Year: 2019

GPS coordinates, latitude and longitude with Google Maps: 37.82,57.69

INPUT DATA

Geometric characteristics of the river basins

 $F = 44.57 \text{ km}^2$ (Surface area of the drainage basin)

O = 47.37 km (Length of the watershed)

 $Fv = 23.2 \text{ km}^2$ (Surface area of greater portion of the drainage basin)

 $Fm = 21.36 \text{ km}^2$ (Surface area of smaller portion of the drainage basin)

Lv = 9.45 km (Natural length of main water course)

Lb = 9.79 km (Length of the drainage basin measured by a series of paraller lines)

Topograpfic characteristics of the river basins

Contour line length - Liz [km]: ["1.22 ","34.72 ","30.98 ","23 ","22.07 ","3.11 "]

The area between the two neighboring contour lines - f [km²]: ["0.13 ","11.60 ","12.29 ","6.51 ","11.62 ","2.15 ","0.27 "]

h0 = 1300 m (Altitude of the initial contour)

Ah = 100 m (Equidistance)

Hmin = 1291 (Lowest altitude in the drainage basin)

Hmax = 1874 (Highest altitude in the draigane basin

Hydrological characteristics of the river basins

 $\Sigma L = 72.2$ km (The total length of the main watercourse with tributaries of 1st and 2nd class)

Lm = 9.2 km (The shortest distance between the fountain (head and mouth))

Water permeability

fp = 0.31 (Part of the surface area of the drainage basin which is composed of highly water permeable structures from the rocks (limestone, sand, gravel))

fpp = 0.16 (Part of the surface area of the drainage basin which is composed of the rocks of medium water permeability (schist, marls, sandstone))

fo = 0.53 (Part of the surface area of the drainage basin which is composed of the rocks of poor water permeability (heavy clay, compact eruptive))

Land use

fs = 0 (Part of the surface area of the drainage basin under the forest)

ft = 0.97810 (Part of the surface area of the drainage basin which is under the grass, meadows, pastures and orchards)

fg = 0.02190 (Part of the surface area of the drainage basin which is bare or under the soils without grass vegetation)

Meteorological data

hb = 34.89 mm (Level of torrent rain)

Up (years) = 100

to = 10.80 °C (Average annual air temperature)

Hgod = 319.6 mm (Average annual quantity of precipitation)

Erosion coefficients

Y = 0.96577 (Types of soil structures and allied types)

0 % (Sand, gravel and incoherent soils)

0 % (Saline soils)

24.21 % (Decomposed limestone and marls)

44.09 % (Serpentines, red sand stones, flishe deposits)

0 % (Podzols and parapodzols, decomposed schist)

0 % (Solid and Schist limestone, Terra Rosa and Humic soil)

0 % (Brown forest soils and Mountain soils)

```
31.71 % (Epieugleysol and Marshlands)
0 % (Good structured Chernozems and alluvial well-structured deposits)
0 % (Bare, compact igneous)
Xa = 0.61367 (Planning of the drainage basin, rate of drainage basin regulation)
0 % (Bare lands)
2.19 % (Plough-lands)
7.1 % (Orchards and vineyards)
90.71 % (Mountain pastures)
0 % (Meadows)
0 % (Degraded forests)
0 % (Well-constituted forests)
\phi = 0.59227 (Numerical coefficient of visible and clearly pointed processes of soil erosion)
0 % (Depth erosion)
30.79 % (80% of the river basin under rill and gully erosion)
21.5 % (50% of the river basin under rill and gully erosion)
0 % (100% of the river basin under surface erosion)
0 % (100% of the river basin under surface erosion, without visible furrows, ravines and land
slides)
0 % (50% of the river basin under surface erosion)
47.72 % (20% of the river basin under surface erosion)
0 % (There are smaller slides in the watercourse beds)
0 % (The river basin mostly under plough-land)
0 % (The river basin under forests and perennial vegetation)
INPUT DATA
```

A = 0.97747619047619 (Coefficient of the river basin form) m = 0.39930574575061 (Coefficient of the watershed development) B = 4.5526046986721 km (Average river basin width) a = 0.082585278276481 ((A)symmetry of the river basin)

G = 1.6199237155037 (Density of the river network of the basin)

K = 1.0271739130435 (Coefficient of the river basin tortuousness)

 $H_{sr} = 1507.0878393538 \text{ m}$ (Average river basin altitude)

D = 216.0878393538 m (Average elevation difference of the river basin) $I_{sr} = 25.824545658515 \% \text{ (Average river basin decline)}$ $H_{leb} = 583 \text{ m (The height of the local erosion base of the river basin)}$ $E_r = 71.822087476017 \text{ (Coefficient of the erosion energy of the river basins relief)}$ $S_1 = 0.766 \text{ (Coefficient of the regions permeability)}$ $S_2 = 0.80438 \text{ (Coefficient of the vegetation cover)}$ W = 0.46439268369981 m (Analytical presentation of the water retention in inflow) $2gDF^{1/2} = 434.69633849387 \text{ m km s}^{-1} \text{ (Energetic potential of water flow during torrent rains)}$ $Q_{max} = 121.58152098868 \text{ m}^3 \text{ s}^{-1} \text{ (Maximal outflow from the river basin)}$

T = 1.08627804912 (Temperature coefficient of the region)

Z = 0.65219633340214 (Coefficient of the river basin erosion)

 $W_{god} = 25603.999523126 \text{ m}^3 \text{ god}^{-1}$ (Production of erosion material in the river basin

 $R_u = 0.32898587042325$ (Coefficient of the deposit retention)

 $G_{god} = 8423.3540694323 \text{ m}^3 \text{ god}^{-1} \text{ (Real soil losses)}$

 $G_{god} \text{ km}^{-2} = 188.99156539 \text{ m}^3 \text{ km}^{-2} \text{ god}^{-1} \text{ (Real soil losses per km}^2\text{)}$

http://www.wintero.me