Web application for Intensity of Erosion and Outflow

Name of the River Basin: Shirindareh S2-intB

Country: Iran, Islamic Republic of

Year: 2019

GPS coordinates, latitude and longitude with Google Maps: 37.77,57.63

INPUT DATA

Geometric characteristics of the river basins

 $F = 10.86 \text{ km}^2$ (Surface area of the drainage basin)

O = 17.64 km (Length of the watershed)

 $Fv = 7.75 \text{ km}^2$ (Surface area of greater portion of the drainage basin)

 $Fm = 3.1 \text{ km}^2$ (Surface area of smaller portion of the drainage basin)

Lv = 3.3 km (Natural length of main water course)

Lb = 6.39 km (Length of the drainage basin measured by a series of paraller lines)

Topograpfic characteristics of the river basins

Contour line length - Liz [km]: ["10.70 ","16.54 ","2.57 "] The area between the two neighboring contour lines - f [km²]: ["3.17 ","6.56 ","1.12 ","0.01 "] h0 = 1300 m (Altitude of the initial contour)

 $\Delta h = 100 \text{ m (Equidistance)}$

Hmin = 1251 (Lowest altitude in the drainage basin)

Hmax = 1565 (Highest altitude in the draigane basin

Hydrological characteristics of the river basins

 $\Sigma L = 18.79$ km (The total length of the main watercourse with tributaries of 1st and 2nd class)

Lm = 3.17 km (The shortest distance between the fountain (head and mouth))

Water permeability

fp = 0.3 (Part of the surface area of the drainage basin which is composed of highly water permeable structures from the rocks (limestone, sand, gravel))

fpp = 0.53 (Part of the surface area of the drainage basin which is composed of the rocks of medium water permeability (schist, marls, sandstone))

fo = 0.17 (Part of the surface area of the drainage basin which is composed of the rocks of poor water permeability (heavy clay, compact eruptive))

Land use

fs = 0 (Part of the surface area of the drainage basin under the forest)

ft = 1.00000 (Part of the surface area of the drainage basin which is under the grass, meadows, pastures and orchards)

fg = 0.00000 (Part of the surface area of the drainage basin which is bare or under the soils without grass vegetation)

Meteorological data

hb = 33.67 mm (Level of torrent rain)

Up (years) = 100

to = 11.70 °C (Average annual air temperature)

Hgod = 304.9 mm (Average annual quantity of precipitation)

Erosion coefficients

Y = 0.95085 (Types of soil structures and allied types)

0 % (Sand, gravel and incoherent soils)

0 % (Saline soils)

0 % (Decomposed limestone and marls)

70.17 % (Serpentines, red sand stones, flishe deposits)

0 % (Podzols and parapodzols, decomposed schist)

0 % (Solid and Schist limestone, Terra Rosa and Humic soil)

0 % (Brown forest soils and Mountain soils)

```
29.83 % (Epieugleysol and Marshlands)
0 % (Good structured Chernozems and alluvial well-structured deposits)
0 % (Bare, compact igneous)
Xa = 0.6209 (Planning of the drainage basin, rate of drainage basin regulation)
0 % (Bare lands)
0 % (Plough-lands)
20.9 % (Orchards and vineyards)
79.1 % (Mountain pastures)
0 % (Meadows)
0 % (Degraded forests)
0 % (Well-constituted forests)
\phi = 0.6574 (Numerical coefficient of visible and clearly pointed processes of soil erosion)
0 % (Depth erosion)
9.13 % (80% of the river basin under rill and gully erosion)
54.53 % (50% of the river basin under rill and gully erosion)
0 % (100% of the river basin under surface erosion)
9.99 % (100% of the river basin under surface erosion, without visible furrows, ravines and
land slides)
0 % (50% of the river basin under surface erosion)
26.35 % (20% of the river basin under surface erosion)
0 % (There are smaller slides in the watercourse beds)
0 % (The river basin mostly under plough-land)
0 % (The river basin under forests and perennial vegetation)
INPUT DATA
```

m = 0.28248415675116 (Coefficient of the watershed development)

B = 1.6995305164319 km (Average river basin width)

a = 0.85714285714286 ((A)symmetry of the river basin)

G = 1.7302025782689 (Density of the river network of the basin)

K = 1.0410094637224 (Coefficient of the river basin tortuousness)

 $H_{sr} = 1338.7348066298 \text{ m}$ (Average river basin altitude)

D = 87.7348066298 m (Average elevation difference of the river basin) $I_{sr} = 27.449355432781 \% \text{ (Average river basin decline)}$ $H_{leb} = 314 \text{ m (The height of the local erosion base of the river basin)}$ $E_r = 55.058239303942 \text{ (Coefficient of the erosion energy of the river basins relief)}$ $S_1 = 0.661 \text{ (Coefficient of the regions permeability)}$ $S_2 = 0.8 \text{ (Coefficient of the vegetation cover)}$ W = 0.46175990646396 m (Analytical presentation of the water retention in inflow) $2gDF^{1/2} = 136.72576933407 \text{ m km s}^{-1} \text{ (Energetic potential of water flow during torrent rains)}}$ $Q_{max} = 34.799843908111 \text{ m}^3 \text{ s}^{-1} \text{ (Maximal outflow from the river basin)}}$ T = 1.1269427669585 (Temperature coefficient of the region) $Z = 0.69743174994669 \text{ (Coefficient of the river basin erosion)}}$ $W_{god} = 6827.9692143301 \text{ m}^3 \text{ god}^{-1} \text{ (Production of erosion material in the river basin)}}$

 $G_{god} = 1277.3360254956 \text{ m}^3 \text{ god}^{-1} \text{ (Real soil losses)}$

 $G_{god} \text{ km}^{-2} = 117.61841855392 \text{ m}^3 \text{ km}^{-2} \text{ god}^{-1} \text{ (Real soil losses per km}^2\text{)}$

 $R_u = 0.18707407508733$ (Coefficient of the deposit retention)

http://www.wintero.me